

DOUGLAS ALGEBRAS WITH NO MAXIMAL SUBALGEBRA AND NO MINIMAL SUPERALGEBRA

CARROLL GUILLORY

ABSTRACT. We give several examples of Douglas Algebras that do not have any maximal subalgebra. We find a condition on these algebras that guarantees that some do not have any minimal superalgebra. We also show that if A is the only maximal subalgebra of a Douglas algebra B , then the algebra A does not have any maximal subalgebra.

1. INTRODUCTION

Let $\mathbf{D}$ denote the open unit disk in the complex plane and $\mathbf{T}$ the unit circle. By L^∞ we mean the space of essentially bounded measurable functions on $\mathbf{T}$ with respect to the normalized Lebesgue measure. We denote by H^∞ the space of all bounded analytic functions in $\mathbf{D}$. Via identification with boundary functions, H^∞ can be considered as a uniformly closed subalgebra of L^∞ . Any uniformly closed subalgebra B strictly between L^∞ and H^∞ is called a Douglas algebra. $M(B)$ will denote the maximal ideal space of a Douglas algebra B . If we let $X = M(L^\infty)$, we can identify L^∞ with $C(X)$, the algebra of continuous functions on X . If C is the set of all continuous functions on $\mathbf{T}$, we set

$$H^\infty + C = \{h + g : g \in C, h \in H^\infty\}$$

$H^\infty + C$ then becomes the smallest Douglas algebra containing H^∞ properly. The function

$$q(z) = \prod_{n=1}^{\infty} \frac{|z_n|}{z_n} \frac{z - z_n}{1 - \bar{z}_n z}$$

is called a Blaschke product if $\sum_{n=1}^{\infty} (1 - |z_n|)$ converges. The set $\{z_n\}$ is called the zero set of q in $\mathbf{D}$. Here $\frac{|z_n|}{z_n} = 1$ is understood whenever $z_n = 0$. We call q an interpolating Blaschke product if

$$\inf_n \prod_{m \neq n} \left| \frac{z_m - z_n}{1 - \bar{z}_n z_m} \right| > 0.$$

Research at MSRI is partially supported by NSF grant DMS-9022140.

An interpolating Blaschke product q is called sparse (or thin) if

$$\lim_{n \rightarrow \infty} \prod_{m \neq n} \left| \frac{z_m - z_n}{1 - \bar{z}_n z_m} \right| = 1.$$

The set

$$Z(q) = \{x \in M(H^\infty) \setminus \mathbf{D} : \mathbf{q}(\mathbf{x}) = \mathbf{0}\}$$

is called the zero set of b in $M(H^\infty + C)$. Any function h in H^∞ with $|h| = 1$ almost everywhere on $\mathbf{T}$ is called an inner function. Since $|q| = 1$ for any Blaschke product, Blaschke products are inner functions. Let

$$QC = (H^\infty + C) \cap \overline{(H^\infty + C)}$$

and, for $x \in M(H^\infty + C)$, set

$$Q_x = \{y \in M(L^\infty) : f(x) = f(y) \text{ for all } f \in QC\}.$$

Q_x is called the QC-level set for x . For $x \in M(H^\infty + C)$, we denote by u_x the representing measure for x and its support set by $\text{supp } u_x$. By $H^\infty[\bar{q}]$ we mean the Douglas algebra generated by H^∞ and the complex conjugate of the function q . Since X is the Shilov boundary for every Douglas algebra, a closed set E contained in X is called a peak set for a Douglas algebra B if there is a function in B with $f = 1$ on E and $|f| < 1$ on $X \setminus E$. A closed set E is a weak peak set for B if E is the intersection of a family of peak sets. If the set E is a weak peak set for H^∞ and we define

$$H_E^\infty = \{f \in L^\infty : f|_E \in H^\infty|_E\},$$

then H_E^∞ is a Douglas algebra. For a Douglas algebra B , B_E is similarly defined. A closed set E contained in X is called the essential set for B , denoted by $\text{ess}(B)$, if E is the smallest set in X with the property that for f in L^∞ with $f = 0$ on E , then f is in B .

For an interpolating Blaschke product q we put $N(\bar{q})$ the closure of

$$\bigcup \{\text{supp } u_x : x \in M(H^\infty + C) \text{ and } |q(x)| < 1\}.$$

$N(\bar{q})$ is a weak peak set for H^∞ and is referred to as the nonanalytic points of q . By $N_0(\bar{q})$ we denote the closure of

$$\bigcup \{\text{supp } u_x : x \in Z(q)\}.$$

For an $x \in M(H^\infty)$ we let

$$E_x = \{y \in M(H^\infty) : \text{supp } u_y = \text{supp } u_x\}$$

and call E_x the level set of x . Since the sets $\text{supp } u_x$ and $N(\bar{q})$ are weak peak sets for H^∞ , both $H_{\text{supp } u_x}^\infty$ and $H_{N(\bar{q})}^\infty$ are Douglas algebras. For any interpolating Blaschke

product q we set

$$A = \bigcap_{x \in M(H^\infty + C) \setminus M(H^\infty[\bar{q}])} H_{\text{supp } u_x}^\infty$$

and

$$A_0 = \bigcap_{y \in Z(q)} H_{\text{supp } u_y}^\infty.$$

It is easy to see that $A \subseteq A_0$ and it was shown in [11] that $A = H_{N(q)}^\infty$. For x and y in $M(H^\infty)$, the pseudo-hyperbolic distance is defined by

$$\rho(x, y) = \sup \{|h(x)| : |h| \leq 1, h \in H^\infty, h(y) = 0\}.$$

For x and y in $\mathbf{D}$ we have

$$\rho(x, y) = \left| \frac{x - y}{1 - \bar{y}x} \right|.$$

For any $x \in M(H^\infty)$, we define the Gleason part of x by

$$P_x = \{y \in M(H^\infty) : \rho(x, y) < 1\}.$$

If $P_x \neq \{x\}$, then x is said to be a nontrivial point. We denote by G the set of nontrivial points of $M(H^\infty + C)$, and for a Douglas algebra B , we set

$$G_B = G \cap (M(H^\infty + C) \setminus M(B)).$$

A point x in G_B is called a minimal support point of G_B (or simply a minimal support point of B) if there are no $y \in G_B$ such that $\text{supp } u_y \subseteq \text{supp } u_x$. The set $\text{supp } u_x$ is called a minimal support set for B . For Douglas algebras B and B_0 with $B_0 \subseteq B$ we let $\Omega(B, B_0)$ be all interpolating Blaschke products q such that $\bar{q} \in B$ but $\bar{q} \notin B_0$.

We denote by $\Omega(B)$ the set of all interpolating Blaschke products q with $\bar{q} \in B$. Let B be a Douglas algebra. The Bourgain algebra B_b of B relative to L^∞ is the set of those elements of L^∞ , f , such that $\|ff_n + B\|_\infty \rightarrow 0$ for every sequence $\{f_n\}$ in B with $f_n \rightarrow 0$ weakly. The minimal envelop B_m of a Douglas algebra B is defined to be the smallest Douglas algebra which contains all minimal superalgebra of B . An algebra A is called a minimal superalgebra of B if for all $x, y \in M(B) \setminus M(A)$, $x \neq y$ implies $\text{supp } u_x = \text{supp } u_y$.

A major part of this paper was done while the author was at the Mathematical Sciences Research Institute. The author thanks the Institute for its support and also thanks Pamela Gorkin for the many valuable discussions.

2. A CONDITION FOR DOUGLAS ALGEBRAS TO HAVE EQUAL ESSENTIAL SETS

Consider the Douglas algebras A and A_0 defined above. In [11] some conditions were given when $A \subsetneq A_0$ but yet $\text{ess}(A) = \text{ess}(A_0)$. This happened because $\text{ess}(A) = N(\bar{q})$ and $\text{ess}(A_0) = N_0(\bar{q})$ (this is not hard to show). Theorem 1 of [11] gives conditions when $\text{ess}(A) \neq \text{ess}(A_0)$. The conditions found in Theorem 5 of [11] are

far more complicated than those found in Theorem 1 below. Yet $\text{ess}(A) = \text{ess}(A_0)$ in that Theorem (Theorem 5 of [11]) and also satisfies the condition in Theorem 1.

Theorem 1. *Let B_0 be a subalgebra of a Douglas algebra B with*

$$\text{ess}(B_0) \neq X.$$

If for every $x \in M(B_0) \setminus M(B)$ we have $\text{ess}(H_{\text{supp } u_x}^\infty) = \text{ess}(B_{\text{supp } u_x})$, then $\text{ess}(B) = \text{ess}(B_0)$.

Proof: We note that $\text{ess}(H_{\text{supp } u_x}^\infty) = \text{supp } u_x$. Hence if $\text{ess}(H_{\text{supp } u_x}^\infty)$ is contained in $\text{ess}(B)$ for every $x \in M(B_0) \setminus M(B)$, then $\text{supp } u_y \subset \text{ess}(B)$ for every $y \in M(B_0)$ and so we get $\text{ess}(B_0) \subset \text{ess}(B)$. Since $B_0 \subset B$ we have that $\text{ess}(B) \subset \text{ess}(B_0)$, and we get $\text{ess}(B) = \text{ess}(B_0)$.

Corollary 1. *Let B_0 be a maximal subalgebra of a Douglas algebra B . Then $\text{ess}(B_0) = \text{ess}(B)$.*

Since $M(B_0) = M(B) \cup E_x$ for some $x \in M(B_0) \setminus M(B)$, we have that if z and y are in $M(B_0) \setminus M(B)$, then $\text{supp } u_y = \text{supp } u_x = \text{supp } u_z$. Now $\text{ess}(B_{\text{supp } u_x}) = \text{ess}(H_{\text{supp } u_x}^\infty)$ since the set

$$\bigcup \{ \text{supp } u_y : y \in M(B) \cap M(H_{\text{supp } u_x}^\infty) \}$$

is dense in $\text{supp } u_x$ (because x is a minimal support point of B). Thus $\text{ess}(B) = \text{ess}(B_0)$.

Corollary 2. *Let A be any Douglas algebra and q be an interpolating Blaschke product with $\bar{q} \notin A$. Then $\text{ess}(A) = \text{ess}(A[\bar{q}])$.*

Proof: Let $B = A[\bar{q}]$ and let $x \in M(A) \setminus M(B)$. We need the following two facts:

- i) $M(A) = \{x \in M(H^\infty + C) : A_{\text{supp } u_x} = H_{\text{supp } u_x}^\infty\}$
and
- ii) If $A \subset B$, then $A_{\text{supp } u_x} \subset B_{\text{supp } u_x}$.

We use i) and ii) to show that $B_{\text{supp } u_x} = A_{\text{supp } u_x}[\bar{q}]$. Since $A \subset B$ by ii) $M(B_{\text{supp } u_x}) \subset M(A_{\text{supp } u_x})$. By the Chang–Marshall Theorem [1,16] it suffices to show that $M(B_{\text{supp } u_x}) = M(A_{\text{supp } u_x}[\bar{q}])$. Let $y \in M(B_{\text{supp } u_x})$. Since $M(B_{\text{supp } u_x}) \subset M(A_{\text{supp } u_x})$ we have that $y \in M(A_{\text{supp } u_x})$ and $|q(y)| = 1$ (since $M(B) = \{y \in M(A) : |q(y)| = 1\}$). Hence $y \in M(A_{\text{supp } u_x}[\bar{q}])$. Therefore $M(B_{\text{supp } u_x}) \subset M(A_{\text{supp } u_x}[\bar{q}])$.

Now suppose $y \notin M(B_{\text{supp } u_x})$. If $y \notin M(A_{\text{supp } u_x})$, then $y \notin A_{\text{supp } u_x}[\bar{q}]$ and we have nothing to prove. So we can assume that $y \in M(A_{\text{supp } u_x})$. Now since $y \notin M(B_{\text{supp } u_x})$ implies that $|q(y)| < 1$ and $y \in M(A_{\text{supp } u_x})$. Hence $y \notin M(A_{\text{supp } u_x}[\bar{q}])$. Thus $M(A_{\text{supp } u_x}[\bar{q}]) \subset M(B_{\text{supp } u_x})$. By the Chang–Marshall Theorem we have that $M(A_{\text{supp } u_x}[\bar{q}]) = M(B_{\text{supp } u_x})$. Hence $A_{\text{supp } u_x}[\bar{q}] = B_{\text{supp } u_x}$. Thus we have

$$\begin{aligned} \text{ess}(B_{\text{supp } u_x}) &= \text{ess}(A_{\text{supp } u_x}[\bar{q}]) \\ &= \text{ess}(H_{\text{supp } u_x}^\infty[\bar{q}]) \quad \text{by i)} \end{aligned}$$

So it suffices to show that

$$\text{ess} (H_{\text{supp } u_x}^\infty [\bar{q}]) = \text{ess} (H_{\text{supp } u_x}^\infty).$$

To do this set $B_1 = H_{\text{supp } u_x}^\infty$ and

$$E = \bigcup \{ \text{supp } u_y : \text{supp } u_y \subset \text{supp } u_x, |q(y)| = 1 \}.$$

Assume that $\bar{E}$, the closure of E , is properly contained in $\text{supp } u_x$. Put

$$B_2 = \bar{H}_E^\infty = \overline{\{f \in L^\infty : f|_E \in H^\infty|_E\}}.$$

By [2, page 39] $M(B_2) = \{m \in M(H^\infty) : \text{supp } u_m \subseteq \bar{E}\} \cup M(L^\infty)$. Since $\bar{E} \subsetneq \text{supp } u_x$ we have that $B_1 \subsetneq B_2$. Therefore $M(B_2) \subsetneq M(B_1)$ and so there is a nontrivial point $x_0 \in M(B_1) \setminus M(B_2)$ [5, Proposition 4.1] such that (a) $\text{supp } u_{x_0} \subsetneq \text{supp } u_x$, (b) $\text{supp } u_{x_0} \not\subseteq \bar{E}$ [otherwise $x_0 \in M(B_2)$], (c) $|q(x_0)| < 1$ and (d) $\text{supp } u_{x_0} \cap E = \emptyset$. In (a) $\text{supp } u_{x_0} \subsetneq \text{supp } u_x$ for if $\text{supp } u_{x_0} = \text{supp } u_x$ for all $x_0 \in M(B_1) \setminus M(B_2)$, then B_1 is a maximal subalgebra of B_2 , so by Corollary 1 we have $\bar{E} = \text{supp } u_x = \text{ess}(B_1)$ (in fact B_2 has no maximal subalgebras). The fact that $|q(x_0)| < 1$ implies that $\bar{q} \notin H_{\text{supp } u_{x_0}}^\infty$.

Therefore there is a $y_0 \in Z(q)$ such that $\text{supp } u_{y_0} \subseteq \text{supp } u_{x_0}$. By Theorem 2 of [10] there is a $z_0 \in Z(q)$ such that $\text{supp } u_{z_0}$ is a minimal support set for $H^\infty[\bar{q}]$ that is contained in $\text{supp } u_{x_0} \subseteq \text{supp } u_x$. Since q is an interpolating Blaschke product $\text{supp } u_{z_0}$ is not trivial. By [5, Theorem 4.2] there is $m \in M(H^\infty + C)$ so that $\text{supp } u_m$ is nontrivial and $\text{supp } u_m \subsetneq \text{supp } u_{z_0} \subseteq \text{supp } u_{x_0}$. Since $\text{supp } u_{z_0}$ is a minimal support set for $H^\infty[\bar{q}]$ we have that $|q(m)| = 1$. Thus $\text{supp } u_{x_0} \cap E \neq \emptyset$. This contradicts (d). So $\bar{E} = \text{supp } u_x$ and $\text{ess}(H_{\text{supp } u_x}^\infty) = \text{ess}(H_{\text{supp } u_x}^\infty[\bar{q}])$ and we get $\text{ess}(B_{\text{supp } u_x}) = \text{ess}(H_{\text{supp } u_x}^\infty)$ for every $x \in M(A) \setminus M(B)$, so Corollary 2 follows from our Theorem.

We mention here that Corollary 2 was proved in [20, Theorem 2] by another method.

There are algebras B_0 and B that satisfy the hypothesis of Theorem 1 and are not of the form $B = B_0[\bar{q}]$ for any interpolating Blaschke product (if $B_0 \subseteq B$). To see this let Γ be the collection of sparse Blaschke products and B be the Douglas algebra $[H^\infty : \bar{q}; q \in \Gamma]$. Let q_0 be any element in Γ and put $B_0 = H^\infty[\bar{q}]$. Then $B_0 \subset B$. By a Theorem of Hedenmalm's [13] we have that if b is a Blaschke product such that $\bar{b} \in B$, then $b = b_1 \cdots b_n$ where each b_i , $i = 1, \dots, n$, is a sparse Blaschke product. Hence if $x \in M(B_0) \setminus M(B)$, then x is the zero of some sparse Blaschke product. So x is a minimal support point of B for every $x \in M(B_0) \setminus M(B)$. This implies that $\text{ess}(B_{\text{supp } u_x}) = \text{ess}(H_{\text{supp } u_x}^\infty)$. So, by Theorem 1, we have $\text{ess}(B) = \text{ess}(B_0)$ (Theorem 2 below shows that $H_{\text{supp } u_x}^\infty$ is a maximal subalgebra of $B_{\text{supp } u_x}$.) Now suppose there is a Blaschke product $q \in \Omega(B, B_0)$ with $B = B_0[\bar{q}]$. Again by Hedenmalm's Theorem we have $q = q_1 \cdots q_n$ with each q_i a sparse Blaschke product. Let Q be any infinite sparse Blaschke product such that $|Q| = 1$ on $\bigcup_{x \in Z(q)} P_x$. Then there is a $m \in M(H^\infty + C)$ such that $Q(m) = 0$ but $m \notin \bigcup_{x \in Z(q)} P_x$. Thus $|q(m)| = 1$ and so we get that

$m \in M(B_0[\bar{q}])$. Thus $\bar{Q} \notin B_0[\bar{q}]$ and yet $\bar{Q} \in B$. This implies that $B_0[\bar{q}] \subsetneq B$, which is a contradiction.

3. MAXIMAL SUBALGEBRAS THAT HAVE NO MAXIMAL SUBALGEBRA

We begin by extending Proposition 1 of [9]. There the authors showed that if $x \in Z(q)$ with q a sparse Blaschke product, then the algebra $H_{\text{supp } u_x}^\infty$ is a maximal subalgebra of $H_{\text{supp } u_x}^\infty[\bar{q}]$. Below we show that this is true for a larger class.

Theorem 2. *Let A be any Douglas algebra with maximal subalgebra and x be a minimal support point of G_A . Then $H_{\text{supp } u_x}^\infty$ is a maximal subalgebra of $A_{\text{supp } u_x}$ and $A_{\text{supp } u_x} = H_{\text{supp } u_x}^\infty[\bar{q}]$ for some $q \in \Omega(A)$.*

Proof: Let $B_0 = H_{\text{supp } u_x}^\infty$ and $B = A_{\text{supp } u_x}$. Suppose x is a minimal support set for G_A . Then we have for any interpolating Blaschke product $\psi \in \Omega(A)$ with $|\psi(x)| < 1$ and any $y \in M(H^\infty + C)$ with $\text{supp } u_y \subsetneq \text{supp } u_x$, $|\psi(y)| = 1$. Thus if $\psi_0 \in \Omega(B)$ and $|\psi_0(x)| < 1$ there is a $\psi \in \Omega(A)$ such that $\psi|_{\text{supp } u_x} = \psi_0|_{\text{supp } u_x}$. This implies that $|\psi_0(y)| = 1$ for every such y . Hence x is a minimal support point for G_B . Note that this implies that $M(B) = M(B_0) \setminus E_x$ where E_x is the level set for x . Hence $M(B_0) = M(B) \cup E_x$, so by Theorem 1 of [10], B_0 is a maximal subalgebra of B . Let q be any element in $\Omega(A)$ with $q(x) = 0$. Then $q \in \Omega(B, B_0)$ and we have that $B = B_0[\bar{q}]$.

Theorem 3. *A Douglas algebra A has no maximal subalgebra if and only if $H_{\text{supp } u_x}^\infty$ is not a maximal subalgebra of $A_{\text{supp } u_x}$ for every $x \in G_A$.*

Proof: Suppose A has no maximal subalgebra and let $x \in G_A$. Since x is not minimal support set of G_A there is a $y \in G_A$ with $\text{supp } u_y \subsetneq \text{supp } u_x$, and a $\psi \in \Omega(A)$ such that $|\psi(y)| < 1$. Since $\bar{\psi} \notin H_{\text{supp } u_y}^\infty$, we can assume that $\psi(y) = 0$. Hence $y \notin M(A_{\text{supp } u_x})$. By Lemma 4 of [8] there is a $\psi_0 \in \Omega(A_{\text{supp } u_x})$ such that $|\psi_0(y)| = 1$ and $\psi_0(x) = 0$. Then we have $H_{\text{supp } u_x}^\infty \subsetneq H_{\text{supp } u_x}^\infty[\bar{\psi}_0] \subsetneq A_{\text{supp } u_x}$. So $H_{\text{supp } u_x}^\infty$ is not a maximal subalgebra of $A_{\text{supp } u_x}$.

Suppose that for all $x \in G_A$, $H_{\text{supp } u_x}^\infty$ is not a maximal subalgebra of $A_{\text{supp } u_x}$. Then there is an algebra B with $H_{\text{supp } u_x}^\infty \subsetneq B \subseteq A_{\text{supp } u_x}$. Thus we can find a $y \in M(H^\infty + C)$ such that $\text{supp } u_y \subsetneq \text{supp } u_x$ and $y \in M(B) \setminus M(A_{\text{supp } u_x})$. This implies that there is an interpolating Blaschke product q with $\bar{q} \in B \subset A_{\text{supp } u_x}$ such that $|q(y)| = 1$ and $|q(x)| < 1$. Hence there is a $q_0 \in \Omega(A)$ with $q_0|_{\text{supp } u_x} = \bar{q}|_{\text{supp } u_x}$. So $|q_0(y)| = 1$ and $|q_0(x)| < 1$. This implies that x is not a minimal support point of G_A for every $x \in G_A$. So by Theorem 1 of [10] A has no maximal subalgebra.

Proposition 1. *Let $x \in M(H^\infty) \setminus M(L^\infty)$. Then $H_{\text{supp } u_x}^\infty$ has no maximal subalgebra.*

Proof: Now $\text{ess}(H_{\text{supp } u_x}^\infty) = \text{supp } u_x$. Hence if $y \in G_{H_{\text{supp } u_x}^\infty}$, then $\text{supp } u_y \cap \text{supp } u_x = \emptyset$. Hence if A is a subalgebra of $H_{\text{supp } u_x}^\infty$, then there is a $y \in M(A) \setminus M(H_{\text{supp } u_x}^\infty)$ with

$\text{supp } u_y \cap \text{supp } u_x = \emptyset$. Hence $\text{ess}(A) \supseteq \text{supp } u_y \cup \text{supp } u_x \supsetneq \text{supp } u_x = \text{ess}(H_{\text{supp } u_x}^\infty)$. By Corollary 1, A is not a maximal subalgebra of $H_{\text{supp } u_x}^\infty$.

Proposition 2. *Let A be a Douglas algebra that has only one maximal subalgebra A_0 . Then A_0 has no maximal subalgebra.*

Proof: Suppose there is a subalgebra $B_0 \subseteq A_0$ such that B_0 is a maximal subalgebra of A_0 . Then by Theorem 1 of [10] there is an $x_0 \in G_{A_0}$ such that

$$M(B_0) = M(A_0) \cup E_{x_0} \quad (1)$$

Since A_0 is a maximal subalgebra of A there is an $x \in G_A \cap M(B_0)$ such that

$$M(A_0) = M(A) \cup E_x \quad (2)$$

By (1) and (2) we have that $M(B_0) = M(A) \cup E_x \cup E_{x_0}$. Since $x \in M(A_0)$ we have that $\text{supp } u_x \neq \text{supp } u_{x_0}$. Also since $x_0 \notin E_x$ and $x_0 \in G_{A_0}$ implies that $\text{supp } u_{x_0} \not\subseteq \text{supp } u_x$ (otherwise $x_0 \in M(A)$ by (2)). We show that x_0 is a minimal support point of G_A , and hence get a contradiction. Let $y \in M(H^\infty + C)$ such that $\text{supp } u_y \subsetneq \text{supp } u_{x_0}$. Since x_0 is a minimal support point of G_{A_0} we have that $y \in M(A_0) = M(A) \cup E_x$. If $y \in M(A)$ then we are done. So we can assume $y \in E_x$. If $y \in E_x$, then $\text{supp } u_y = \text{supp } u_x$, so we have that $\text{supp } u_x \subsetneq \text{supp } u_{x_0}$. Since $x \notin M(A)$ there is an interpolating Blaschke product q with $\bar{q} \in A$ and such that $q(x) = 0$. By Theorem 2 of [11] there is an uncountable set U of $Z(q)$ such that (a) $\text{supp } u_m \subsetneq \text{supp } u_{x_0}$ for all $m \in U$ and (b) $\text{supp } u_m \cap \text{supp } u_k$ for all $m, k \in U, m \neq k$. By (1) each such $m \in U$ is in $M(A_0)$. Since for all $m \in U$ (except if $m = x$) we have $\text{supp } u_x \cap \text{supp } u_m = \emptyset$, hence by (2), $m \in M(A)$. But $\bar{q} \in A$ and $U \subset Z(q) \cap M(A)$. This is a contradiction, and we get $y \notin E_x$. So $y \in M(A)$ and since $\text{supp } u_{x_0} \neq \text{supp } u_x$ we have that x_0 is a minimal support point of G_A . This is a contradiction. So A_0 has no maximal subalgebra.

Note that Proposition 1 follows from Proposition 2 if x is a minimal support point for some interpolating Blaschke product.

Let q be an interpolating Blaschke product. We consider the algebra $H_{N(\bar{q})}^\infty$. Certainly $H_{N(\bar{q})}^\infty$ is not known to be a maximal subalgebra of any Douglas algebra, but does have some of the same properties of $H_{\text{supp } u_x}^\infty$. For example

Proposition 3. *The algebra $H_{N(\bar{q})}^\infty$ has no maximal subalgebra.*

Proof: Set $B = H_{N(\bar{q})}^\infty$. Let $x \in G_B$ and suppose x is a minimal support point for G_B . Then if $y \in M(H^\infty + C)$ such that $\text{supp } u_y \subsetneq \text{supp } u_x$ we have that $y \in M(B)$. By [2, page 39] we must have that $\text{supp } u_y \subseteq N(\bar{q}) = \text{ess}(B)$. Thus we have that $\text{ess}(B) \cap \text{supp } u_x = N(\bar{q}) \cap \text{supp } u_x \neq \emptyset$. By [14, Theorem 1] $N(\bar{q}) = \bigcup_{x \in Z(q)} Q_x$. So there is an $x_0 \in Z(q)$ such that $\text{supp } u_x \cap Q_{x_0} \neq \emptyset$. By the definition of Q_{x_0} , we have that $\text{supp } u_x \subset Q_{x_0}$. By [2, page 39] this implies that $x \in M(B)$, which is a contradiction. So if $x \in G_B$, then $\text{supp } u_x \cap N(\bar{q}) = \emptyset$, which implies that x is not a minimal support point for G_B . B has no maximal subalgebra.

4. MINIMAL SUPERALGEBRAS OF $H_{\text{supp } u_x}^\infty$

We will compute the Bourgain algebras and the Minimal Envelopes of the Douglas algebra $H_{\text{supp } u_x}^\infty$ for any $x \in M(H^\infty + C)$. We have the following

Theorem 4. *Let $x \in M(H^\infty + C) \setminus M(L^\infty)$ such that $|q(x)| < 1$ for some interpolating Blaschke product q and set $B = H_{\text{supp } u_x}^\infty$. Then*

- i) *Either $B_b = B$ or $B_b = B[\bar{\psi}]$ for some interpolating Blaschke product ψ .*
- ii) *Either $B_m = B_b = B$ or $B_m = B[\bar{\psi}]$ for some interpolating Blaschke product ψ .*

Proof: We will use Theorem 2 of [4] which says that for any interpolating Blaschke product ψ with $\bar{\psi} \in B_b$, we have that the set $Z(\psi) \cap M(B)$ is a finite set and the fact that

$$M(B) = M(L^\infty) \cup \{m \in M(H^\infty) : \text{supp } u_m \subseteq \text{supp } u_x\} \quad (3)$$

We claim that if ψ is an interpolating Blaschke product such that $\bar{\psi} \in B_b$, then $Z(\psi) \cap M(B) \subset E_x$, the level set of x . Suppose not. Then there is an $x_0 \in Z(\psi) \cap M(B)$ such that $\text{supp } u_{x_0} \subsetneq \text{supp } u_x$. By Theorem 2 of [11] there is an uncountable set Γ of $Z(\psi)$ such that (a) $\text{supp } u_\gamma \subsetneq \text{supp } u_x$ for all $\gamma \in \Gamma$ and (b) $\text{supp } u_{x_m} \cap \text{supp } u_\gamma = \emptyset$ for all $m, \gamma \in \Gamma$, $m \neq \gamma$. By (a) and (3) each $\gamma \in M(B)$ and so $\Gamma \subset Z(\psi) \cap M(B)$. This implies that the set $Z(\psi) \cap M(B)$ is infinite. This is a contradiction. Hence if $x_0 \in Z(\psi) \cap M(B)$ then $\text{supp } u_{x_0} = \text{supp } u_x$, so we get $Z(\psi) \cap M(B) \subset E_x$. There are two possibilities (1) the set $Z(\psi) \cap M(B) = \emptyset$ for which $\bar{\psi} \in B$, so $B_b \subseteq B$. This gives us the case when $B_b = B$. (2) If $Z(\psi) \cap M(B) \neq \emptyset$ but finite. Then the algebra $B[\bar{\psi}] \subseteq B_b$. To show that $B_b = B[\bar{\psi}]$, let ψ_0 be other interpolating Blaschke product with $\bar{\psi}_0 \in B_b$. Since both sets $Z(\psi_0) \cap M(B)$ and $Z(\psi) \cap M(B)$ are contained in E_x , we have that $M(B) \setminus M(B[\bar{\psi}_0]) = M(B) \setminus M(B[\bar{\psi}]) = E_x$. Thus $M(B[\bar{\psi}_0]) = M(B[\bar{\psi}])$ and by the Chang–Marshall Theorem [1, 16] we have $B[\bar{\psi}_0] = B[\bar{\psi}]$, since this is true for all ψ, ψ_0 we have by Theorem C of [12], $B_b = B[\bar{\psi}]$ for any such ψ $B[\bar{\psi}]$ is a minimal superalgebra of B .) This proves (i).

To prove (ii) let $\bar{\psi} \in B_m$. Then, by Theorem 3 of [12] there is a finite set $\{x_1, \dots, x_n\} \subset Z(\psi) \cap M(B)$ such that $\{u \in M(B) : |\psi(u)| < 1\} = E_{x_1} \cup \dots \cup E_{x_n}$. Again we claim that $E_{x_1} = E_{x_2} = \dots = E_{x_n} = E_x$. Suppose that $E_{x_1} \neq E_{x_2}$. Then $\text{supp } u_{x_1} \neq \text{supp } u_{x_2}$. By (3) either $\text{supp } u_{x_1} \subsetneq \text{supp } u_x$ or $\text{supp } u_{x_2} \subsetneq \text{supp } u_x$ or both. Let us suppose that $\text{supp } u_{x_1} \subsetneq \text{supp } u_x$. Then by Theorem 2 of [11] there is an uncountable set Γ such that $E_\alpha \neq E_\beta$ for all $\alpha, \beta \in \Gamma$ and $\bigcup_{\alpha \in \Gamma} E_\alpha \subset \{u \in M(B) : |\psi(u)| < 1\}$. This contradicts Theorem 3 of [12]. Thus $E_{x_1} = E_{x_2} = \dots = E_{x_n} = E_x$. As before we have that for $\bar{\psi} \in B_m$, $Z(\psi) \cap M(B) \subset E_x$ and $B_m = B[\bar{\psi}]$ if $Z(\psi) \cap M(B) \neq \emptyset$. This proves (ii).

Corollary 3. i) *Let $x \in M(H^\infty + C) \setminus M(L^\infty)$ and $B = H_{\text{supp } u_x}^\infty$. Then $B \subset B_m$ if and only if x is a minimal support point of $H^\infty[\bar{\psi}]$ for some interpolating Blaschke product ψ .*

- ii) $B = B_b = B_m$ if and only if x is not a minimal support point of $H^\infty[\bar{\psi}]$ for any interpolating Blaschke product.

Theorem 4 i) has also appeared in [17].

REFERENCES

- [1] S. Y. Chang, *A characterization of Douglas Subalgebras*, Acta. Math. **137** (1976), 81–89.
- [2] T. Gamelin, *Uniform Algebras*, Prentice-Hall, Englewood Cliffs, NJ, 1969.
- [3] J. Garnett, *Bounded Analytic Functions*, Academic Press, New York, 1981.
- [4] P. Gorkin, K. Izuchi, R. Mortini, *Bourgain Algebras of Douglas Algebras*, Canad. J. Math. **44** (1992), 797–804.
- [5] P. Gorkin, R. Mortini, *Interpolating Blaschke Products and Factorization in Douglas Algebras*, Michigan Math. J. **38** (1991), 147–160.
- [6] C. J. Guillory, *Douglas Algebras of the Form $H^\infty[\bar{q}]$* , J. Math. Anal. Appl. **126** (1987), 269–274.
- [7] C. J. Guillory, *A Characterization of a Sparse Blaschke Product*, Canad. Math. Bull. **32** (1989), 385–390.
- [8] C. J. Guillory, *A Note on Exposed Points and Restricted Douglas Algebras*, to be submitted.
- [9] C. J. Guillory, K. Izuchi, D. Sarason, *Interpolating Blaschke Products and Divisors in Douglas Algebras*, Proc. Royal Irish Acad. **18** (1984), 1–7.
- [10] C. J. Guillory, K. Izuchi, *Maximal Douglas Subalgebras and Minimal Support Sets*, Proc. Amer. Math. Soc. **116** (1992), 477–481.
- [11] C. J. Guillory, K. Izuchi, *Interpolating Blaschke Products and Nonanalytic Sets*, J. Complex Variables **23** (1993), 163–175.
- [12] C. J. Guillory, K. Izuchi, *Minimal Envelopes of Douglas Algebras and Bourgain Algebras* Houston J. Math. Vol. **19** (1993), 201–222.
- [13] H. Hedenmalm, *Thin Interpolating Sequences and Three Algebras of Analytic Functions*, Proc. Amer. Math. Soc. **99** (1987), 489–495.
- [14] K. Izuchi, *QC-Level Sets and Quotients of Douglas Algebras*, J. Funct. Anal. **65** (1986), 293–308.
- [15] K. Izuchi, *Countably Generated Douglas Algebras*, Trans. Amer. Math. Soc. **299**, No 3(1986), 171–192.
- [16] D. Marshall, *Subalgebras of L^∞ Containing H^∞* , Acta Math. **137** (1976), 91–98.
- [17] R. Mortini, R. Younis, *Douglas Algebras Which are Invariant under the Bourgain Map*, Arch. Math. **59** (1992), 371–378.
- [18] R. Younis, *M-Ideals of L^∞/H^∞ and Support Sets*, Illinois J. Math. **29**, **1** (1985), 96–102.
- [19] R. Younis, *Division in Douglas Algebras and Some Applications*, Arch. Math. **45** (1985), 550–560.
- [20] R. Younis, *Best Approximation in Certain Douglas Algebras*, Proc. Amer. Math. Soc. **80** (1980), 639–642.

UNIVERSITY OF SOUTHWESTERN LOUISIANA, LAFAYETTE, LOUISIANA 70504
 Current address: MSRI, 1000 Centennial Drive, Berkeley, CA 94720
 E-mail address: cjg@usl.edu